

Information geometry of the spherical model

W. Janke

Institut für Theoretische Physik, Universität Leipzig, Augustusplatz 10/11, D-04109 Leipzig, Germany

D. A. Johnston

Department of Mathematics, Heriot-Watt University, Riccarton, Edinburgh, EH14 4AS, Scotland

R. Kenna

School of Mathematical and Information Sciences, Coventry University, Coventry, CV1 5FB, England

(Received 28 October 2002; published 14 April 2003)

Motivated by the observation that geometrizing statistical mechanics offers an interesting alternative to more standard approaches, we calculate the scaling behavior of the curvature \mathcal{R} of the information geometry metric for the spherical model. We find that $\mathcal{R} \sim \epsilon^{-2}$, where $\epsilon = \beta_c - \beta$ is the distance from criticality. The discrepancy from the naively expected scaling $\mathcal{R} \sim \epsilon^{-3}$ is explained and compared with that for the Ising model on planar random graphs, which shares the same critical exponents.

DOI: 10.1103/PhysRevE.67.046106

PACS number(s): 05.50.+q, 75.10.Hk

I. THE INFORMATION GEOMETRY OF SPIN MODELS

The idea of endowing the space of parameters with a metric and geometrical structure has been borrowed from parametric statistics [1] and employed to some effect in statistical mechanics [2–8]. The approach seems to be particularly fruitful for a spin model in a field where the parameters are β , the inverse temperature, and h , the external field. In this case the (Fisher-Rao) metric is simply given by

$$G_{ij} = \partial_i \partial_j f, \quad (1)$$

where f is the reduced free energy per site and $\partial_i = (\partial/\partial\beta, \partial/\partial h)$.

For such a metric the scalar curvature may be calculated as

$$\mathcal{R} = -\frac{1}{2G^2} \begin{vmatrix} \partial_\beta^2 f & \partial_\beta \partial_h f & \partial_h^2 f \\ \partial_\beta^3 f & \partial_\beta^2 \partial_h f & \partial_\beta \partial_h^2 f \\ \partial_\beta^2 \partial_h f & \partial_\beta \partial_h^2 f & \partial_h^3 f \end{vmatrix}, \quad (2)$$

where $G = \det(G_{ij})$ is the determinant of the metric itself.

The work in Refs. [2–5] has made it clear that, as one might expect, the scalar curvature plays a central role in any attempt to look at statistical mechanics from a geometrical perspective. In particular, for all the models that have been considered so far, the curvature diverges only at a phase transition point for physical ranges of the parameter values. We assume a standard scaling form for the free energy per site,

$$f(\epsilon, h) = \lambda^{-1} f(\epsilon \lambda^{a_\epsilon}, h \lambda^{a_h}), \quad (3)$$

where $\epsilon = \beta_c - \beta$ (with β_c marking the critical point), and a_ϵ and a_h are the scaling dimensions for the energy and spin operators, respectively.

The spherical model has a one-sided critical point [9] and we are interested in the high-temperature domain (where $\epsilon > 0$). There, standard scaling assumptions allow us to write Eq. (3) as

$$f(\epsilon, h) = \epsilon^{1/a_\epsilon} \psi_+(h \epsilon^{-a_h/a_\epsilon}), \quad (4)$$

where we have introduced the scaling function ψ_+ . Consideration of the behavior of the components of \mathcal{R} in Eq. (2) then leads to the scaling

$$\mathcal{R} \sim \epsilon^{\alpha-2}, \quad (5)$$

for the curvature itself.

Equation (5) shows that the scalar curvature not only characterizes phase transition points as divergences, as do the more standard statistical mechanical quantities such as the specific heat C and susceptibility χ , it displays a scaling behavior that allows the extraction of the critical exponent α . Indeed, with hyperscaling, Eq. (5) can be recast as

$$\mathcal{R} \sim \xi^d, \quad (6)$$

where ξ is the correlation length and d is the dimensionality of the system. So the curvature is proportional to the correlation volume, an intuitively reasonable result on dimensional grounds.

At first sight, our calculation of the scaling behavior of \mathcal{R} in Ref. [10] for the Ising model on planar random graphs puts a spanner in the works. There, $\alpha = -1$, but $\mathcal{R} \sim \epsilon^{-2}$ rather than the expected $\mathcal{R} \sim \epsilon^{-3}$. However, returning to the detailed scaling of the individual terms in Eq. (2) showed that a negative α affected some of the scaling of the components and the end result could be traced back to these modifications. In principle this modified behavior should apply to *any* model with negative α , so an interesting test would be to calculate \mathcal{R} , or at least its scaling limit, for another model with this property.

The spherical model provides just such a test case. It was solved (in field) in the classic Berlin and Kac paper [9] and the critical exponents in three dimensions found to be $\alpha=-1$, $\beta=1/2$, $\gamma=2$, which are identical to those of the Ising model on the two-dimensional (2D) planar random graphs [11]. This is remarkable, because there are no obvious physical similarities between the two models. An additional motivation for any such calculation is the general paucity of spin models that have been solved in field. Indeed, \mathcal{R} for spin models has been obtained explicitly only for the 1D Ising [3], Bethe lattice Ising [6], and 1D Potts [12] models and the scaling form calculated for the Ising model on planar random graphs [10]. This forms a rather small sample on which to formulate hypotheses about its general behavior and its scaling properties. In the sequel, we extend this short list incrementally by looking at the scaling behavior of \mathcal{R} in the spherical model. We concentrate on the 3D case, but also discuss other dimensions, including the mean-field-like behavior that sets in at $d=4$.

II. THE SPHERICAL MODEL

Berlin and Kac [9] introduced the spherical model (and the Gaussian model) in an attempt to understand how generic some of the features of Onsager's solution [13] of the 2D Ising model are for ferromagnetic spin models, particularly for other dimensions. In the spherical model, the ± 1 condition on the value of the Ising spins is relaxed, whilst retaining a global constraint on the *total* spin magnitude. With s_i denoting the value of a spin at a site i of a hypercubic lattice, the partition function is [9]

$$\mathcal{Z} = \int ds_1 \cdots ds_N \exp\left(\beta \sum_{\langle ij \rangle} s_i s_j + h \sum_i s_i\right) \delta\left(\sum_i s_i^2 - N\right), \quad (7)$$

where N is the total number of sites. This can be evaluated by exponentiating the constraint and using steepest descent, resulting in the following expression for the reduced free energy per site in the thermodynamic limit, $N \rightarrow \infty$:

$$f = \frac{1}{2} \log\left(\frac{\pi}{\beta}\right) + \beta z - \frac{1}{2} g(z) + \frac{h^2}{4\beta(z-d)}, \quad (8)$$

where

$$g(z) = \frac{1}{(2\pi)^d} \int_0^{2\pi} d\omega_1 \cdots d\omega_d \log\left(z - \sum_{k=1}^d \cos(\omega_k)\right). \quad (9)$$

The saddle point value of z , which appears in the expression for the free energy in Eq. (8), is determined from

$$g'(z) = 2\beta - \frac{h^2}{2\beta(z-d)^2}. \quad (10)$$

The solution reveals no transition for $d=1$ and 2, and a transition with the exponents $\alpha=-1$, $\beta=1/2$, $\gamma=2$ for $d=3$. For $d \geq 4$, on the other hand, mean-field behavior with $\alpha=0$, $\beta=1/2$, $\gamma=1$ sets in (modified by multiplicative logarithmic corrections in the $d=4$ case [14]). For the following formulas we confine our attention to the $d=3$ case.

It is useful to note that, with $h=0$, Eq. (10) gives

$$\frac{dz}{d\beta} = \frac{2}{g''(z)}, \quad (11)$$

and hence

$$\frac{d^2z}{d\beta^2} = -\frac{4g^{(3)}(z)}{[g''(z)]^3}. \quad (12)$$

The critical point is given by $z=d=3$ and $h=0$ [9], and the behavior of $g(z)$ in this region is determined by differentiating Eq. (9) twice and then expanding for the small ω_k values that give the dominant contribution. One finds

$$g''(z) \sim -\frac{1}{2\sqrt{2}\pi}(z-3)^{-1/2}. \quad (13)$$

A further differentiation gives

$$g^{(3)}(z) \sim \frac{1}{4\sqrt{2}\pi}(z-3)^{-3/2}, \quad (14)$$

and an integration yields

$$g'(z) = \frac{1}{\sqrt{2}\pi}(z-3)^{1/2} + g'(3), \quad (15)$$

where $g'(3) = (18 + 12\sqrt{2} - 10\sqrt{3} - 7\sqrt{6})[K(2\sqrt{3} + \sqrt{6} - 2\sqrt{2} - 3)]^2 \approx 0.505462019 \cdots$ is the exactly known massless 3D lattice propagator at the origin, with $K(k^2)$ denoting the standard elliptic integral. This latter expression can be combined with Eq. (10) with $h=0$ to give

$$(z-3) \sim 8\pi^2(\beta_c - \beta)^2 \sim \epsilon^2, \quad (16)$$

in which $\beta_c = g'(3)/2 \approx 0.252731009 \dots$. Equations (13) and (14) may then be substituted in Eqs. (11) and (12) to give the scaling of $dz/d\beta$ and $d^2z/d\beta^2$,

$$\lim_{z \rightarrow 3} \frac{dz}{d\beta} = \lim_{z \rightarrow 3} [-4\sqrt{2}\pi(z-3)^{1/2}] = 0,$$

$$\lim_{z \rightarrow 3} \frac{d^2z}{d\beta^2} = 16\pi^2, \quad (17)$$

which we shall employ below in the calculation of the scalar curvature.

III. THE SCALAR CURVATURE

We recapitulate the general considerations of Ref. [10] before going on to discuss in detail the results for the spheri-

cal model. In Eq. (4), we define $A = 1/a_\epsilon$ and $C = -a_h/a_\epsilon$, which, in terms of the standard exponents, gives $A = 2 - \alpha$ and $A + C = \beta$. If $A > 2$ (which is the case for $\alpha < 0$), the specific heat, rather than diverging, will be a constant at the critical point, which we denote by $A(A - 1)\phi(0)$. With this in mind, substitution of the scaling function into Eq. (2) gives

$$\mathcal{R} = -\frac{1}{2G^2} \begin{vmatrix} A(A-1)\phi(0) & 0 & \epsilon^{A+2C}\psi_+''(0) \\ -A(A-1)(A-2)\epsilon^{A-3}\psi_+(0) & 0 & -(A+2C)\epsilon^{A+2C-1}\psi_+''(0) \\ 0 & -(A+2C)\epsilon^{A+2C-1}\psi_+''(0) & 0 \end{vmatrix}. \quad (18)$$

In Eq. (18) the terms with an odd number of h derivatives have been set to zero. These do not appear because of the $h \rightarrow -h$ symmetry of the free energy. The scaling for the determinant of the metric G is given by

$$G = A(A-1)\epsilon^{A+2C}\phi(0)\psi_+''(0), \quad (19)$$

so the leading term in \mathcal{R} (for $A > 2$) is

$$\mathcal{R} = \frac{(A+2C)^2}{2A(A-1)\phi(0)}\epsilon^{-2}, \quad (20)$$

or, translating back to the standard critical exponents,

$$\mathcal{R} = \frac{\gamma^2}{2(2-\alpha)(1-\alpha)\phi(0)}\epsilon^{-2}. \quad (21)$$

In summary, if $\alpha < 0$, the expected scaling of \mathcal{R} is $\mathcal{R} \sim \epsilon^{-2}$ rather than $\mathcal{R} \sim \epsilon^{\alpha-2}$, which is seen for positive α .

We now move on to examine the scaling of the various terms contributing to \mathcal{R} in Eq. (2) for the spherical model itself. As we have remarked, the $h \rightarrow -h$ symmetry in the free energy per site, f , of the spherical model means that any terms with an odd number of h derivatives will automatically be zero when $h=0$, hence $f_{\beta h} = f_{\beta\beta h} = f_{hhh} = 0$. This leaves the nonzero terms (again when $h=0$)

$$\begin{aligned} f_{\beta\beta} &= \frac{\partial z}{\partial \beta} + \frac{1}{2\beta^2}, \\ f_{hh} &= \frac{1}{2\beta(z-3)}, \\ f_{\beta\beta\beta} &= \frac{\partial^2 z}{\partial \beta^2} - \frac{1}{\beta^3}, \\ f_{hh\beta} &= -\frac{1}{2\beta(z-3)^2} \frac{\partial z}{\partial \beta} - \frac{1}{2\beta^2(z-3)}, \end{aligned} \quad (22)$$

which, using Eqs. (16) and (17), have the following behavior in the scaling region:

$$\begin{aligned} f_{\beta\beta} &\sim \frac{1}{2\beta_c^2}, \\ f_{hh} &\sim \frac{1}{16\pi^2\beta_c(\beta_c-\beta)^2} \sim \epsilon^{-2}, \\ f_{\beta\beta\beta} &\sim 16\pi^2 - \frac{1}{\beta_c^3}, \\ f_{hh\beta} &\sim \frac{1}{8\pi^2\beta_c(\beta_c-\beta)^3} - \frac{1}{16\pi^2\beta_c^2(\beta_c-\beta)^2} \sim \epsilon^{-3}. \end{aligned} \quad (23)$$

Comparing with the individual terms in Eq. (18) we see that the expected general scaling of each term (for $\alpha < 0$) does indeed apply and that overall we have, as in Eq. (21),

$$\mathcal{R} \sim \epsilon^{-2}. \quad (24)$$

We thus see that calculating the scaling of \mathcal{R} for the 3D spherical model for which $\alpha = -1$ gives results in accordance with expectations from general scaling arguments, which take into account the negative α , similar to the Ising model on planar random graphs.

In $d \geq 4$ dimensions the exponents of the spherical model attain their mean-field values, $\alpha=0$, $\beta=1/2$, $\gamma=1$. In this case $g'(z) \sim z - d + g'(d)$ near criticality and the resulting scaling of the components of the expression for \mathcal{R} is

$$\mathcal{R} = -\frac{1}{2G^2} \begin{vmatrix} \text{const} & 0 & \epsilon^{-1} \\ \text{const} & 0 & \epsilon^{-2} \\ 0 & \epsilon^{-2} & 0 \end{vmatrix}, \quad (25)$$

where the determinant of the metric scales as $G = \epsilon^{-1}$. Thus, \mathcal{R} itself scales as ϵ^{-2} , in the mean-field case. This is in agreement with the expected scaling of $\epsilon^{\alpha-2}$ when $\alpha \geq 0$ since, in that case [4],

$$\mathcal{R} = -\frac{1}{2G^2} \begin{vmatrix} A(A-1)\epsilon^{A-2}\psi_+(0) & 0 & \epsilon^{A+2C}\psi_+''(0) \\ -A(A-1)(A-2)\epsilon^{A-3}\psi_+(0) & 0 & -(A+2C)\epsilon^{A+2C-1}\psi_+''(0) \\ 0 & -(A+2C)\epsilon^{A+2C-1}\psi_+''(0) & 0 \end{vmatrix}, \quad (26)$$

where the scaling of the metric determinant is

$$G = A(A-1)\epsilon^{2A+2C-2}\psi_+(0)\psi_+''(0). \quad (27)$$

The difference in scaling between the $\alpha \geq 0$ and the $\alpha < 0$ cases originates in the $f_{\beta\beta}$ term which contributes to both \mathcal{R} and G in Eqs. (26) and (27). In the negative α case, the contribution of this term comes from the regular part of the free energy and is nondiverging.

Note that for $\alpha=0$, $A=2$, and although the displayed scaling term $A(A-1)(A-2)\epsilon^{A-3}\psi_+(0)$ in Eq. (26) would be expected to vanish, a regular term could still contribute. This is indeed what is seen in the explicit calculation for the spherical model. Two further caveats arise. First, for the upper critical dimension $d=4$ itself, there are multiplicative logarithmic corrections [14], which require a careful handling. Second, for $d>4$, where the scaling is described by the mean-field theory, the hyperscaling relation fails. So one should be quite cautious in comparing the explicit calculation in terms of ϵ with the expressions involving the correlation length ξ .

IV. CONCLUSIONS

We have calculated the scaling behavior of the scalar curvature \mathcal{R} of the information geometry metric for the 3D

spherical model, finding, as for the Ising model on planar random graphs, that $\mathcal{R} \sim \epsilon^{-2}$. Careful considerations of the scaling of the various elements that contribute to \mathcal{R} show that this is in accordance with the expectations. We also briefly discussed the scaling behavior of \mathcal{R} for $d \geq 4$, corresponding to the mean-field exponents.

The calculation reported here provides another example of a statistical mechanical model in which the curvature of the information geometry metric diverges at the critical point, with a clearly quantifiable scaling behavior. Once again it is curious that the relevant exponent is shared with the Ising model on planar random graphs. All critical exponents now coincide in the two models though there is no obvious physical relation between them. It seems that the suggestion in the original Berlin and Kac paper [9] that the 3D spherical model might provide a hint to the behavior of the 3D Ising model applies rather to the Ising model on 2D planar random graphs.

ACKNOWLEDGMENTS

W.J. and D.A.J. were partially supported by EC IHP network ‘‘Discrete Random Geometries: From Solid State Physics to Quantum Gravity’’ HPRN-CT-1999-000161. D.A.J. and R.K. were also partially supported by an Enterprise Ireland/British Council Research Visits Scheme grant.

-
- [1] R.A. Fisher, *Philos. Trans. R. Soc. London, Ser. A* **222**, 309 (1922); C.R. Rao, *Bull. Calcutta Math. Soc.* **37**, 81 (1945).
 - [2] G. Ruppeiner, *Rev. Mod. Phys.* **67**, 605 (1995); *Phys. Rev. A* **20**, 1608 (1979); **24**, 488 (1980).
 - [3] H. Janyszek, *Rep. Math. Phys.* **24**, 1 (1986); **24**, 11 (1986); H. Janyszek and R. Mrugała, *Phys. Rev. A* **39**, 6515 (1989).
 - [4] H. Janyszek, *J. Phys. A* **23**, 477 (1990).
 - [5] E.J. Brody, *Phys. Rev. Lett.* **58**, 179 (1987); D. Brody and N. Rivier, *Phys. Rev. E* **51**, 1006 (1995); D. Brody and A. Ritz, *Nucl. Phys. B* **522**, 588 (1998); D. Brody and L. Hughston, *Proc. R. Soc. London, Ser. A* **455**, 1683 (1999).
 - [6] B.P. Dolan, *Proc. R. Soc. London, Ser. A* **454**, 2655 (1998).
 - [7] B.P. Dolan, *Int. J. Mod. Phys. A* **12**, 2413 (1997).
 - [8] D. O’Connor and C.R. Stephens, *Prog. Theor. Phys.* **90**, 747 (1993); B.P. Dolan, *Int. J. Mod. Phys. A* **10**, 2439 (1995); **10**, 2703 (1995); *Nucl. Phys. B* **528**, 553 (1998); J. Gaiete and D. O’Connor, *Phys. Rev. D* **54**, 5163 (1996); A.C. Irving and J.C. Sexton, *ibid.* **55**, 5456 (1997).
 - [9] T. Berlin and M. Kac, *Phys. Rev.* **86**, 821 (1952).
 - [10] W. Janke, D.A. Johnston, and Ransinghe P.K.C. Malmini, *Phys. Rev. E* **66**, 056119 (2002).
 - [11] V.A. Kazakov, *Phys. Lett. A* **119**, 140 (1986); D.V. Boulatov and V.A. Kazakov, *Phys. Lett. B* **186**, 379 (1987).
 - [12] B.P. Dolan, D.A. Johnston, and R. Kenna, *J. Phys. A* **35**, 9025 (2002).
 - [13] L. Onsager, *Phys. Rev.* **65**, 117 (1944).
 - [14] R. Kenna and C.B. Lang, *Phys. Lett. B* **264**, 396 (1991); *Nucl. Phys. B* **393**, 461 (1993).